

LETTER TO THE EDITOR

Tentative detection of phosphine in IRC +10216 [★]

M. Agúndez¹, J. Cernicharo¹, J. R. Pardo¹, M. Guélin², T. G. Phillips³

¹ Departamento de Astrofísica Molecular e Infrarroja, Instituto de Estructura de la Materia, CSIC, Serrano 121, 28006 Madrid, Spain; e-mail: marce@damir.iem.csic.es, cerni@damir.iem.csic.es, pardo@damir.iem.csic.es

² Institut de Radioastronomie Millimétrique, 300 rue de la Piscine, 38406 St. Martin d'Hères, and LERMA/École Normale Supérieure, 24 rue Lhomond, 75231 Paris, France; e-mail: guelin@iram.fr

³ California Institute of Technology, Downs Laboratory of Physics 320-47, Pasadena, CA 91125, USA; e-mail: tgp@submm.caltech.edu

Received ; accepted

ABSTRACT

Aims. The $J_K = 1_0-0_0$ rotational transition of phosphine (PH_3) at 267 GHz has been tentatively identified with a $T_{\text{MB}} \sim 40$ mK spectral line observed with the IRAM 30-m telescope in the C-star envelope IRC +10216.

Methods. A radiative transfer model has been used to fit the observed line profile.

Results. The derived PH_3 abundance relative to H_2 is 6×10^{-9} , although it may have a large uncertainty due to the lack of knowledge about the spatial distribution of this species. If our identification is correct, it implies that PH_3 has a similar abundance to that reported for HCP in this source, and that these two molecules (HCP and PH_3) together take up about 5 % of phosphorus in IRC +10216. The abundance of PH_3 , as that of other hydrides in this source, is not well explained by conventional gas phase LTE and non-LTE chemical models, and may imply formation on grain surfaces.

Key words. Stars: individual: IRC +10216 — stars: carbon — radio lines: stars — astrochemistry — line:identification — stars: AGB and post-AGB

1. Introduction

Of the nearly 150 molecules detected so far in interstellar and circumstellar media, around 3/4 can be formed from just four elements (H, C, N, and O). The remaining 1/4 contain metals (Na, K, Al, Mg, and Fe), halogens (F and Cl), and to a large extent the second-row elements Si, P, and S. The scarcity of molecules containing the second-row elements Si, P, and S, compared to their first-row analogues C, N, and O, reflects on one hand a lower cosmic abundance (Si/C $\sim 1/8$, P/N $\sim 1/300$, and S/O $\sim 1/30$; Asplund et al. 2005) and on the other hand important chemical differences. Second-row elements have generally a larger refractory character, i.e. they tend to form solid condensates and deplete from the gas phase (Field, 1974; Turner, 1991; Lodders & Fegley, 1999). Also, chemical bonds formed by first-row elements are generally stronger and the resulting molecules are more stable than the corresponding second-row analogues. For example, the high stability of C-C bonds allows to form a large variety of organic molecules, while a Si-based chemistry is much more limited.

Concerning phosphorus, the very limited number of interstellar and circumstellar P-bearing molecules known for a long time, just PN and CP (Turner & Bally, 1987; Ziurys, 1987; Guélin et al., 1990) in contrast with the nearly 50 N-bearing

molecules found, may have been related to a sort of mixing of all the factors mentioned above. First, the more than a hundred fold decrease in the cosmic abundance of P compared to that of N. Second, the likely strong depletion of phosphorus suggested by observations of some P-bearing species (Turner et al., 1990). And lastly, some important differences between the chemistry of P and that of N, that are indicated by, for example, the non reactivity of PH_n^+ ($n = 0, 3$) ions with H_2 as opposed to the case of NH_n^+ (Thorne et al., 1984; Adams et al., 1990).

In recent years, however, we have seen the discoveries, mostly in circumstellar media, of new phosphorus compounds such as HCP, PO and CCP (Agúndez et al., 2007; Milam et al., 2008; Tenenbaum et al., 2007; Halfen et al., 2008), which permit to discuss the chemistry of phosphorus on a wider observational basis. In this letter we present the tentative detection of PH_3 , the phosphorus analogue of NH_3 , in the carbon-rich circumstellar envelope IRC +10216. This species, known to be abundant in the atmospheres of the giant gaseous planets Jupiter and Saturn (Weisstein & Serabyn, 1996), has never been observed outside the Solar System.

2. Observations and results

The phosphine molecule, PH_3 , is an oblate symmetric top, thus its rotational levels are given by two quantum numbers (J, K), and radiative transitions are only allowed within levels of the same K ladder ($\Delta J = 1, \Delta K = 0$). The K ladders are grouped into two distinct forms: ortho ($K = 3n$, n an integer) and para ($K \neq 3n$) between which both radiative and collisional transitions are severely forbidden. Its rotational spectrum has been extensively investigated in the laboratory, allowing for the very weak "for-

Send offprint requests to: M. Agúndez

[★] Based on observations carried out with the IRAM 30-meter telescope and the Caltech Submillimeter Observatory (CSO). IRAM is supported by INSU/CNRS (France), MPG (Germany) and IGN (Spain). The CSO is operated by the California Institute of Technology under funding from the National Science Foundation, Grant No. AST-0540882.




Fig. 1. Spectra of IRC +10216 observed with the IRAM 30-m telescope showing on top the $J_K = 1_0-0_0$ line of PH₃ blended with a narrower line assigned to the $J = 15-14$ line of SiS $v = 4$. Shown below are the immediate lower rotational transitions of SiS $v = 4$. Fits to the line profiles using the CLASS *shell* method are shown as grey thick lines.

bidden” transitions ($\Delta J = 0$, $\Delta K = \pm 3$) to be measured and for the hyperfine structure due to the ¹H and the ³¹P nuclear spins to be resolved (see Cazoli & Puzzarini 2006 and references therein). In contrast to NH₃, no evidence of inversion doubling has been found in the case of PH₃. The electric dipole moment has been measured as 0.57395 ± 0.0003 D (Davies et al., 1971).

Following the recent detection of HCP in IRC +10216 (Agúndez et al., 2007), it was speculated that PH₃ might be detectable in this source if the PH₃/HCP abundance ratio is similar to the NH₃/HCN one. Prompted by this hypothesis we searched for the fundamental ortho-PH₃ 1_0-0_0 line, at 267 GHz, with the IRAM 30-m telescope. Preliminary observations with a low spectral resolution, 1.25 MHz, were done in 2007 May and we

Table 1. Observed line parameters in IRC +10216

Transition	Cal. Freq. (MHz)	Obs. Freq. (MHz)	v_{exp}^a (km/s)	$\int T_{\text{MB}} dv$ (K km/s)
PH ₃				
$J_K = 1_0-0_0$	266944.514	266944.5(3)	14.1(3)	1.002(40)
SiS $v = 4$				
$J = 15-14$	266941.754	266942.4(7)	4.0(7)	0.237(40)
$J = 14-13$	249155.372	249155.6(4)	3.5(4)	0.172(40)
$J = 13-12$	231366.976	231367.2(3)	2.7(3)	0.256(30)
$J = 12-11$	213576.710	213576.9(5)	3.5(5)	0.207(40)

NOTE.— Number in parentheses are 1σ uncertainties in units of the last digits. Observed frequencies are given in the rest frame of IRC +10216 assuming a systemic velocity of $v_{\text{LSR}} = -26.5$ km s⁻¹.

^a v_{exp} is the half width at zero level.

found significant emission at the frequency of the PH₃ 1_0-0_0 line and with the expected linewidth ($v_{\text{exp}} = 14.5$ km s⁻¹ for most of the molecular lines in IRC +10216; Cernicharo et al. 2000). The line, however, showed a profile unusual for IRC +10216: neither really U-shaped nor flat-topped. Encouraged by this result we returned to the 30-m telescope in 2008 February and April to re-observe this transition with a 4 times higher spectral resolution (320 kHz). Two SIS receivers operating at 1 mm were used simultaneously with upper side band rejections of ~ 10 dB. The local oscillator was shifted by 80 MHz to identify any contribution from the image sideband. An autocorrelator was used as backend to provide the required spectral resolution of 320 kHz. The secondary mirror was wobbed by $\pm 90''$ at a rate of 0.5 Hz. The pointing and focus of the telescope were checked every 1-2 hours on Saturn, which was closer than 10° to IRC +10216. Here we express the intensity scale in units of main beam brightness temperature T_{MB} . The parameter $B_{\text{eff}}/F_{\text{eff}}$, used to convert T_{A}^* into T_{MB} , is 0.51 and the beam size is $9''$ at 267 GHz for the 30-m.

The resulting spectrum at 267 GHz is shown on top of Fig. 1. The line profile indicates the blend of a normal cusped line of width ~ 29 km s⁻¹, characteristic of optically thin lines arising in the outer envelope (e.g. C₂H), with a narrower (~ 6 km s⁻¹) sharply peaked line, characteristic of vibrationally excited lines arising close to the star. The latter line was soon identified as the $J = 15-14$ transition of SiS in its $v = 4$ vibrational state, based on SiS laboratory spectroscopic data (Sanz et al., 2003; Müller et al., 2007). Emission of SiS in vibrationally excited states up to $v = 3$ has been reported toward IRC +10216 (Turner, 1987; Fonfría et al., 2006; Cernicharo et al., 2000). The linewidths of vibrationally excited SiS are unusually narrow implying that the emission arises from the innermost envelope where the gas has not yet reached the terminal expansion velocity of 14.5 km s⁻¹. To constrain the possible contribution of the $J = 15-14$ line of SiS $v = 4$ to the 267 GHz line, we observed the immediate previous J transitions. In Fig. 1 we show the $J = 14-13$ and $J = 13-12$ lines of SiS in the $v = 4$ state (both observed with a spectral resolution of 320 kHz at the time of the 267 GHz observations) and the $J = 12-11$ line (observed in a previous run in 2005 January with a spectral resolution of 1.25 MHz).

Except for the $J = 14-13$ line of SiS $v = 4$, which is partially blended with a fine structure component of the $N = 27-26$ transition of CCC¹³CH, the observations clearly show that SiS $v = 4$ lines are narrow, with expansion velocities of about 3 km s⁻¹ (see Table 1). With this in mind, the observed emission feature at 267 GHz has been fitted by two line components, a narrow one corresponding to the $J = 15-14$ transition of SiS $v = 4$ and a wider line whose width, $v_{\text{exp}} = 14.1 \pm 0.3$ km s⁻¹, agrees

with the expansion velocity of 14.5 km s^{-1} in IRC +10216, and whose center rest frequency, $266944.5 \pm 0.3 \text{ MHz}$, is in very good agreement with the laboratory frequency of the 1_0-0_0 line of PH₃ (see Table 1). The hyperfine structure is not resolved as the components are separated by less than 0.2 km s^{-1} in velocity (Cazzoli & Puzzarini, 2006), which is smaller than the spectral resolution. The good agreement between the observed and laboratory frequencies is the strongest evidence of PH₃ detection in IRC +10216.

Besides the SiS $v = 4$ line, there are some other lines with frequencies close to that of PH₃ 1_0-0_0 . Most of them, such as SO₂ $30_{9,21}-31_{8,24}$ at 266943.344 MHz or CH₃CH₂CN $15_{4,12}-15_{2,13}$ at 266951.639 MHz , are ruled out as likely contributors since many other lines of these species should have been detected. A more plausible species is HC₃N in the $v_7 = 4$ vibrational state, whose transition $J = 29-28 \ell = 0_e$ lies at 266943.313 MHz (Mbosei et al., 2000). Although several lines of HC₃N in the vibrational excited state $v_7 = 1$ have been observed in IRC +10216 (Cernicharo et al., 2000), we rule out that HC₃N $v_7 = 4$ is the main contributor to the 267 GHz emission based on the upper limit of $T_{\text{MB}} < 0.01 \text{ K}$ that we have from our 30-m data archive for the lower- J transition $J = 26-25 \ell = 0_e$ transition at 239370.171 MHz .

We tried to confirm the identification of PH₃ by observing other transitions. The $J = 2-1$ line at 534 GHz is not reachable from the ground due to severe atmospheric absorption. We, thus, searched for the $J = 3-2$ transition at 800 GHz with the Caltech Submillimeter Observatory (CSO). The observations were carried out in 2008 January using the chopping secondary mode with a throw of $\pm 90''$ at a rate of 1.2 Hz . The SIS receiver was tuned in double sideband and a Fast Fourier Transform Spectrometer was used as backend to provide a spectral resolution of 0.12 MHz . The pointing of the telescope was checked on Saturn. The beam size of the CSO at 800 GHz is $11.5''$ and the beam efficiency is 0.28 . In spite of the good atmospheric conditions (zenith sky opacity at 225 GHz was $0.04-0.08$), the high opacity of the atmosphere at 800 GHz (T_{sys} ranged from 3000 K to 8000 K) did not allowed to reach a noise level low enough to confirm or discard the presence of PH₃ (see top panel in Fig. 2).

3. Analysis and discussion

To interpret the observations we have computed line profiles by means of excitation and radiative transfer calculations based in the Large Velocity Gradient (LVG) formalism (Castor, 1970), coupled to the spectral catalog of J. Cernicharo (Cernicharo et al., 2000). We consider separately both the ortho (o-PH₃) and para (p-PH₃) species of phosphine (the ortho-to-para ratio is assumed to be 1, the statistical value). The energy levels and transition frequencies are computed from the rotational constants reported by Cazzoli & Puzzarini (2006). We include rotational levels in the ground vibrational state up to $J_K = 7_6$ for o-PH₃ and up to $J_K = 5_5$ for p-PH₃. As rate coefficients for collisional de-excitation of o-PH₃(p-PH₃) with H₂ and He, we have adopted those computed for collisions of o-NH₃(p-NH₃) with p-H₂ (Danby et al., 1988) and He (Machin & Roueff, 2005) respectively, properly corrected to the case in which inversion doubling is not resolved.

We assume a distance to IRC +10216 of 150 pc and simulate the circumstellar envelope as a spherically distributed gas expanding at a constant velocity of 14.5 km s^{-1} . The gas density and temperature radial profiles are taken from Agúndez & Cernicharo (2006). PH₃ is assumed to be distributed between an inner radius R_{in} and an outer radius R_{out} , with a con-




Fig. 2. The lower panel shows the PH₃ $J_K = 1_0-0_0$ line at 267 GHz as observed with the IRAM 30-m telescope, in which the fit to the SiS $v = 4$ $J = 15-14$ line has been subtracted. The upper panel shows the observation at 800 GHz with the CSO smoothed to a spectral resolution of 2 MHz . The grey thick lines in both panels correspond to the line profiles given by the LVG model.

stant abundance, $x(\text{PH}_3)$, relative to H₂. We take $R_{\text{in}} = 1.3 \times 10^{15} \text{ cm}$ (about 20 stellar radii), the value adopted by Hasegawa et al. (2006) for NH₃. The values of R_{out} and $x(\text{PH}_3)$ are varied until obtaining the best fit to the observed PH₃ 1_0-0_0 line profile. Since both gas density and temperature vary greatly with radius, we have divided the envelope into various shells and solved for the level populations in each shell independently of the others. We then compute the emergent intensity and weight it with the main beam of the selected telescope.

In the bottom panel of Fig. 2 we plot the observed 1_0-0_0 PH₃ line profile, obtained by subtracting the fit to the SiS $v = 4$ blended line. Also shown is the profile resulting from our best LVG model, whose parameters are $R_{\text{out}} = 2 \times 10^{16} \text{ cm}$ and $x(\text{PH}_3) = 6 \times 10^{-9}$. The model indicates that the upper level involved in the 1_0-0_0 PH₃ line is mostly populated by collisions in the inner $4 \times 10^{15} \text{ cm}$, which corresponds to an angular diameter of about $4''$ and is thus spatially diluted in the $9''$ beam of the 30-m telescope. The top panel of Fig. 2 shows the corresponding prediction for the PH₃ $J = 3-2$ line profiles as observed with the CSO. It is seen that the expected intensity is well below the noise of the observed spectrum and thus is consistent with the

IRAM 30-m result for the $J = 1-0$ line. The spatial dilution is also important for the CSO observations and plays against our efforts to detect the $J = 3-2$ lines.

It should be noted that the abundance and distribution of PH₃ derived are very uncertain. Since we have just one single PH₃ line observed, there exists a considerable degeneracy between models with different values of R_{in} , R_{out} and $x(\text{PH}_3)$. Moreover, infrared pumping to excited vibrational states, not considered in our model, may play an important role in the excitation of the rotational levels in the ground vibrational state. PH₃ has indeed many vibrational bands in the spectral region around 10 μm , a wavelength at which the central source has its maximum flux (Cernicharo et al., 1999).

Confirmation of our tentative detection may rely on further observations of other PH₃ transitions. The $J = 2-1$ and $J = 3-2$ lines, at 534 GHz and 800 GHz respectively are observable with the Herschel Space Observatory (HSO), although our LVG model predicts intensities somewhat weaker than with the CSO due to the larger dilution effect. In the case of the Atacama Large Millimeter Array (ALMA) the high angular resolution that it will provide, better than 3'' at 800 GHz, will perfectly fit with the expected size of the $J = 3-2$ emission. The predictions indicate main beam brightness temperatures of about 1 K, which provide a good opportunity of detection in spite of the high atmospheric opacity at this frequency. We note that these predictions are based on our best LVG model, but the $J = 2-1$ and $J = 3-2$ lines could be more intense than expected if infrared pumping is playing an important role, or if PH₃ is present in hotter regions, inner than 1.3×10^{15} cm.

The derived abundance of PH₃, $x(\text{PH}_3) = 6 \times 10^{-9}$, implies that it takes up about 1.3 % of the available phosphorus. The PH₃/HCP ratio is then found to be 1/2.3 (Agúndez et al., 2007), noticeably higher than the NH₃/HCN ratio, that is about 1/50 in IRC +10216 (Hasegawa et al., 2006; Fonfría et al., 2007). The formation of PH₃ in the gas phase is difficult to explain. Its LTE abundance in the inner envelope is rather low, $< 10^{-12}$, and gas phase reactions yield no net formation in the outer envelope (Agúndez et al., 2007). Besides PH₃, other hydrides such as NH₃, CH₄, H₂O, SiH₄, and H₂S are observed in IRC +10216 with relatively high abundances (Keady & Ridgway, 1993; Hasegawa et al., 2006; Agúndez & Cernicharo, 2006). Many of them are usually assumed as parent molecules, i.e. formed in the inner envelope, in most chemical models of IRC +10216 (e.g. MacKay & Charnley 2001; Agúndez & Cernicharo 2006). However, chemical equilibrium calculations, similar to those reported in Agúndez et al. (2007), indicate abundances much lower than observed, except perhaps for CH₄. A widely invoked explanation, when gas phase chemistry fails to explain an observed abundance, is that of grain surface reactions. In the case of hydrides such as PH₃, a likely formation process is the direct hydrogenation of the heavy atom taking place on grain surfaces.

4. Conclusions

We have tentatively detected PH₃ in IRC +10216 through its $J_K = 1_0-0_0$ transition at 267 GHz. The derived abundance relative to H₂ is 6×10^{-9} . Although with a considerable uncertainty, this value is similar to the HCP abundance found in this source (Agúndez et al., 2007). These two species, HCP and PH₃, would then take up about 5 % of the phosphorus in IRC +10216. The formation of PH₃, unlike that of HCP, is difficult to explain in the gas phase and could occur on grain surfaces. It remains a target for the future to confirm this tentative detection by observing the $J = 3-2$ transition, at 800 GHz, with the ALMA facility. Also,

further observations at 267 GHz in other sources such as CRL 2688, where HCP has been also detected (Milam et al., 2008), will be of great interest to support this tentative detection and to understand the chemistry behind it.

Acknowledgements. We thank the IRAM staff, especially C. Thum, for their kindness during the 30-m observations. We also acknowledge funding support from Spanish MEC through grants AYA2006-14876 and ESP2004-665, and from Spanish CAM under PRICIT project S-0505/ESP-0237 (ASTROCAM). MA also acknowledges grant AP2003-4619 from Spanish MEC.

References

- Adams, N. G., McIntosh, B. J., & Smith, D. 1990, *A&A*, 232, 443
- Agúndez, M., & Cernicharo, J., 2006, *ApJ*, 650, 374
- Agúndez, M., Cernicharo, J., & Guélin, M. 2007, *ApJ*, 662, L91
- Asplund, M., Grevesse, N., & Sauval, A., J. 2005, in *ASP Conf. Ser. 336, Cosmic Abundances as Records of Stellar Evolution and Nucleosynthesis*, ed. T. G. Barnes, III & F. N. Bash (San Francisco: ASP), 25
- Castor, J. I. 1970, *MNRAS*, 149, 111
- Cazzoli, G., & Pazzarini, C. 2006, *J. Mol. Spectr.*, 239, 64
- Cernicharo, J., Yamamura, I., González-Alfonso, E., et al. 1999, *ApJ*, 526, L41
- Cernicharo, J., Guélin, M., & Kahane, C. 2000, *A&AS*, 142, 181
- Danby, G., Flower, D. R., Valiron, P., et al. 1988, *MNRAS*, 235, 229
- Davies, P. B., Neumann, R. M., Wofsy, S. C., & Klemperer, W. 1971, *J. Chem. Phys.*, 55, 3564
- Field, G. B. 1974, *ApJ*, 187, 453
- Fonfría, J. P., Agúndez, M., Tercero, B., et al. 2006, *ApJ*, 646, L127
- Fonfría, J. P., Cernicharo, J., Richter, M. J., & Lacy, J. 2007, *ApJ*, 673, 445
- Guélin, M., Cernicharo, J., Paubert, G., & Turner, B. E. 1990, *A&A*, 230, L9
- Halfen, D. T., Clouthier, D. J., & Ziurys, L. M. 2008, *ApJ*, 677, L101
- Hasegawa, T. I., Kwok, S., Koning, N., et al. 2006, *ApJ*, 637, 791
- Keady, J. J., & Ridgway, S. T. 1993, *ApJ*, 406, 199
- Lodders, K., & Fegley, B. Jr. 1999, in *IAU Symp. 191, Asymptotic Giant Branch Stars*, ed. T. Le Bertre, A. Lèbre, & C. Waelkens (San Francisco: ASP), 279
- Machin, L., & Roueff, E. 2005, *J. Phys. B: Mol. Opt. Phys.*, 38, 1519
- MacKay, D. D. S., & Charnley, S. B. 2001, *MNRAS*, 325, 545
- Mbosei, L., Fayt, A., Dréan, P., & Cosléou, J. 2000, *J. Mol. Struct.*, 517, 271
- Milam, S. N., Halfen, D. T., Tenenbaum, E. D., et al. 2008, *ApJ*, in press (preprint doi:10.1086/589135)
- Müller, H. S. P., McCarthy, M. C., Bizzocchi, L., et al. 2007, *Phys. Chem. Chem. Phys.*, 9, 1579
- Sanz, M. E., McCarthy, M. C., & Thaddeus, P. 2003, *J. Chem. Phys.*, 119, 11715
- Tenenbaum, E. D., Woolf, N. J., & Ziurys, L. M. 2007, *ApJ*, 666, L29
- Thorne, L. R., Anicich, V. G., Prasad, S. S., & Huntress, W. T. 1984, *ApJ*, 280, 139
- Turner, B. E., & Bally, J. 1987, *ApJ*, 321, L75
- Turner, B. E. 1987, *A&A*, 183, L23
- Turner, B. E., Tsuji, T., Bally, J., Guélin, M., & Cernicharo, J. 1990, *ApJ*, 365, 569
- Turner, B. E., 1991, *ApJ*, 376, 573
- Weisstein, E. W., & Serabyn, E. 1996, *Icarus*, 123, 23
- Ziurys, L. M. 1987, *ApJ*, 321, L81